

Existence Results for a Class of Semi-Linear Evolution Equations

Eduardo Hernández M.*

Departamento de Matemática, Instituto de Ciências Matemáticas e de Computação, Universidade de São Paulo - Campus de São Carlos, Caixa Postal 668, 13560-970 São Carlos SP, Brazil
E-mail: lalohm@icmcs.sc.usp.br

We prove the existence of regular solutions for a class of quasi-linear evolution equation modeled in the form $\frac{d}{dt}(x(t) + g(t, x(t))) = Ax(t) + f(t, x(t))$, where A is the infinitesimal generator of an analytic semigroup of bounded linear operators defined on a Banach space X and f, g are appropriate continuous functions. April, 2001 ICMC-USP

1. INTRODUCTION

The class of equations considered in this paper have the form

$$\begin{aligned} \frac{d}{dt}(x(t) + g(t, x(t))) &= Ax(t) + f(t, x(t)), & t > 0, \\ x(0) &= x_0. \end{aligned} \tag{1}$$

We consider system (1) as a Cauchy problem on a Banach space X , where A is the infinitesimal generator of an analytic semigroup of bounded linear operators $(T(t))_{t \geq 0}$; $f, g : [0, T] \times \Omega \rightarrow X$ are appropriate continuous functions and Ω is an open subset of X .

The case $g \equiv 0$ has an extensive literature. The books of Pazy [14], Krein [10], Goldstein [5] and the references contained therein, give a good account of the more important results.

Throughout this paper X will be a Banach space provided with norm $\|\cdot\|$ and the operator $A : D(A) \subset X \rightarrow X$ will be the infinitesimal generator of an analytic semigroup of bounded linear operators $(T(t))_{t \geq 0}$ on X . For the theory of strongly continuous semigroups, refer to [14] and [5]. We mention here only some notations and properties essential to our purpose. In particular, it is well known that there exist $\tilde{M} \geq 1$ and a real number w such that

$$\|T(t)\| \leq \tilde{M}e^{wt}, \quad t \geq 0.$$

* Research Partially Supported by CNPq grant 300.889/92-5 and FAPESP grant 97/011323-0, Brazil

In what follows we assume that $\|T(t)\|$ is uniformly bounded by \tilde{M} and that $0 \in \rho(A)$. In this case it is possible to define the fractional power $(-A)^\alpha$, for $0 < \alpha < 1$, as a closed linear operator with domain $D((-A)^\alpha)$. Furthermore, the subspace $D((-A)^\alpha)$ is dense in X and the expression

$$\|x\|_\alpha = \|(-A)^\alpha x\|$$

defines a norm on $D((-A)^\alpha)$. Hereafter we represent by X_α the space $D((-A)^\alpha)$ endowed with the norm $\|\cdot\|_\alpha$. The following properties are well known (see [14]):

LEMMA 1.1. *If the above conditions hold:*

1. Let $0 < \alpha \leq 1$. Then X_α is a Banach space.
2. If $0 < \beta \leq \alpha$ then $X_\alpha \rightarrow X_\beta$ is continuous.
3. For every constant $a > 0$, there exists $C_a > 0$ such that

$$\|(-A)^\alpha T(t)\| \leq \frac{C_a}{t^\alpha}, \quad 0 < t \leq a.$$

4. For every $a > 0$ there exists a positive constant C'_a such that

$$\|(T(t) - I)(-A)^{-\alpha}\| \leq C'_a t^\alpha, \quad 0 < t \leq a.$$

By analogy with the abstract Cauchy problem

$$\dot{u}(t) = Au(t) + h(t)$$

we adopt the following definitions:

DEFINITION 1.1. A function $x \in C([0, r] : X)$ is a mild solution of the abstract Cauchy problem (1) if: $x(0) = x_0$; for each $0 \leq t < r$ the function $AT(t-s)g(s, x(s))$, $s \in [0, t)$, is integrable and

$$\begin{aligned} x(t) = & T(t)(x_0 + g(0, x_0)) - g(t, x(t)) - \int_0^t AT(t-s)g(s, x(s))ds \\ & + \int_0^t T(t-s)f(s, x(s))ds \end{aligned} \tag{3}$$

for every $t \in [0, r)$.

DEFINITION 1.2. A function $x \in C([0, r] : X)$ is a classical solution of the abstract Cauchy problem (1) if: $x(0) = x_0$; $x(t) \in D(A)$ for all $t \in (0, r)$; \dot{x} is continuous on $(0, r)$ and $x(\cdot)$ satisfies equation (1) on $(0, r)$.

DEFINITION 1.3. A function $x \in C([0, r] : X)$ is an S-classical (Semi-classical) solution of the abstract Cauchy problem (1) if: $x(0) = x_0$; $\frac{d}{dt}(x(t) + g(t, x(t)))$ is continuous on $(0, r)$; $x(t) \in D(A)$ for all $t \in (0, r)$ and $x(\cdot)$ satisfies equation (1) on $(0, r)$.

The paper is organized as follows. In section (2) we discuss the existence of S-classical and classical solutions to the initial value problem (1). Our results are based on the properties of analytic semigroups and the ideas used in [14] chapter 5. In section (3), some applications are considered.

Throughout this paper we assume that X is an abstract Banach space. The terminology and notations are those generally used in operator theory. In particular, if X, Y are Banach spaces, we indicate by $\mathcal{L}(X : Y)$ the Banach space of the bounded linear operator of X into Y and we abbreviate to $\mathcal{L}(X)$ whenever $X = Y$. In addition $B_r(x : X)$ will denote the closed ball in space X with center at x and radius r .

For a bounded function $\xi : [0, a] \rightarrow X$ and $0 < t < a$ we will employ the notation

$$\|\xi(\cdot)\|_t = \sup\{\|\xi(s)\| : s \in [0, t]\}.$$

Finally for $x_0 \in X$, we will use the notation $x(\cdot, x_0)$ for the mild solution of equation (1).

2. REGULAR SOLUTIONS

The existence of mild solutions for the abstract Cauchy problem (1) follows from Theorems 2.1 and 2.2 in [7]; for this reason we choose to omit the proofs of the next two results.

THEOREM 2.1. *Let $x_0 \in X$ and assume that the following conditions hold:*

a) *There exist $\beta \in (0, 1)$ and $L \geq 0$ such that the function g is X_β -valued and satisfies the Lipschitz condition*

$$\|(-A)^\beta g(t, x) - (-A)^\beta g(s, y)\| \leq L(|t - s| + \|x - y\|) \quad (4)$$

for every $0 \leq s, t \leq T$ and $x, y \in \Omega$, and

$$\|(-A)^{-\beta}\| L < 1. \quad (5)$$

b) *The function f is continuous and takes bounded sets into bounded sets.*

c) *The semigroup $(T(t))_{t \geq 0}$ is compact.*

Then there exists a mild solution $x(\cdot, x_0)$ of the abstract Cauchy problem (1) defined on $[0, r]$ for some $0 < r < T$.

THEOREM 2.2. *Let $x_0 \in X$ and assume that the following conditions hold:*

a) There exist $\beta \in (0, 1)$ and $L \geq 0$ such that the function g is X_β -valued and satisfies the Lipschitz condition

$$\| (-A)^\beta g(t, x) - (-A)^\beta g(s, y) \| \leq L(|t - s| + \|x - y\|) \quad (6)$$

for every $0 \leq s, t \leq T$ and $x, y \in \Omega$ and

$$\| (-A)^{-\beta} \| L < 1. \quad (7)$$

b) The function f is continuous and there exists $N > 0$ such that

$$\| f(t, x) - f(s, y) \| \leq N(|t - s| + \|x - y\|) \quad (8)$$

for every $0 \leq s, t \leq T$ and $x, y \in \Omega$.

Then there exists a unique mild solution $x(\cdot, x_0)$ of the abstract Cauchy problem (1) defined on $[0, r]$ for some $0 < r \leq T$.

The existence of S-classical and classical solutions, requires some additional assumptions on the functions g, f . In particular, in the next result we assume that the following assumption hold:

Assumption(f, g) : There exist $0 < \alpha < \beta < 1$ and an open set $\Omega_\alpha \subset X_\alpha$ such that the functions $f, (-A)^\beta g$ are continuous on $[0, T] \times \Omega_\alpha$ and

- There exist $L > 0$ and $0 < \gamma_1, \gamma_2 < 1$ such that for every $(t, x_1), (s, x_2) \in [0, T] \times \Omega_\alpha$

$$\| (-A)^\beta g(t, x_1) - (-A)^\beta g(s, x_2) \| \leq L\{|t - s|^{\gamma_1} + \|x_1 - x_2\|_\alpha\},$$

$$\| f(t, x_1) - f(s, x_2) \| \leq N\{|t - s|^{\gamma_2} + \|x_1 - x_2\|_\alpha\},$$

$$L \| (-A)^{\alpha-\beta} \| < 1.$$

THEOREM 2.3. *Let $x_0 \in \Omega_\alpha$ and assume that system (1) satisfies Assumption(f, g), that g is $D(A)$ -valued continuous and that $1 - \beta < \min\{\beta - \alpha, \gamma_1, \gamma_2\}$. Then there exists a unique S-classical solution $x(\cdot, x_0) \in C([0, r] : X)$ for some $0 < r < T$.*

Proof. Let $0 < r_1 < T$ and $\delta > 0$ such that

$$V = \{(t, x) \in [0, r_1] \times X_\alpha : \| (-A)^\alpha x - (-A)^\alpha x_0 \| < \delta\} \subset [0, T] \times \Omega_\alpha.$$

Assuming that the functions $f, (-A)^\beta g$ are bounded on V by $C_1 > 0$, we choice $0 < r < r_1$ such that:

$$\| (T(\cdot) - I)(-A)^\alpha x_0 \|_r \leq \frac{(1 - \mu)\delta}{6}, \quad (9)$$

$$\| (T(\cdot) - I)(-A)^\alpha g(0, x_0) \|_r \leq \frac{(1 - \mu)\delta}{6}, \quad (10)$$

$$\| (-A)^{\alpha - \beta} \| Lr^{\gamma_1} + C_{1 - \beta + \alpha} C_1 \frac{r^{\beta - \alpha}}{\beta - \alpha} + C_\alpha C_1 \frac{r^{1 - \alpha}}{1 - \alpha} < \frac{(1 - \mu)\delta}{6}, \quad (11)$$

$$LC_{1 - \beta + \alpha} \frac{r^{\beta - \alpha}}{\beta - \alpha} + NC_\alpha \frac{r^{1 - \alpha}}{1 - \alpha} < 1 - \mu, \quad (12)$$

where $\mu = \| (-A)^{\alpha - \beta} \| L$ and $C_\alpha, C_{1 - \beta + \alpha}$ are the constants in Lemma 1.1. On the set

$$S = \{y \in C([0, r] : X) : y(0) = (-A)^\alpha x_0, \| y(t) - (-A)^\alpha x_0 \| \leq \delta, t \in [0, r]\}$$

we define the operator

$$\begin{aligned} \Psi(y)(t) &= T(t)(-A)^\alpha(x_0 + g(0, x_0)) - (-A)^\alpha g(t, (-A)^{-\alpha} y(t)) \\ &\quad + \int_0^t (-A)^{1 - \beta + \alpha} T(t - s)(-A)^\beta g(s, (-A)^{-\alpha} y(s)) ds \\ &\quad + \int_0^t (-A)^\alpha T(t - s) f(s, (-A)^{-\alpha} y(s)) ds. \end{aligned}$$

For the mapping Ψ we consider the decomposition $\Psi = \Psi_1 + \Psi_2$, where

$$\begin{aligned} \Psi_1(y)(t) &= T(t)(-A)^\alpha(x_0 + g(0, x_0)) - (-A)^\alpha g(t, (-A)^{-\alpha} y(t)) \\ &\quad + \int_0^t (-A)^{1 - \beta + \alpha} T(t - s)(-A)^\beta g(s, (-A)^{-\alpha} y(s)) ds, \\ \Psi_2(y)(t) &= \int_0^t (-A)^\alpha T(t - s) f(s, (-A)^{-\alpha} y(s)) ds. \end{aligned}$$

Next we will prove that Ψ_1 and Ψ_2 are well defined, that Ψ satisfies a Lipschitz condition and that the ranges of Ψ is contained in S .

Since $T(\cdot)$ is analytic, the function $s \rightarrow AT(t - s)$ is continuous in the uniform operator topology on $[0, t]$, consequently the function $AT(t - s)g(s, (-A)^{-\alpha} y(s))$ is continuous on $[0, t]$. Moreover from lemma (1.1) we have:

$$\| (-A)^{1 - \beta + \alpha} T(t - s)(-A)^\beta g(s, (-A)^{-\alpha} y(s)) \| \leq \frac{C_{1 - \beta + \alpha} C_1}{(t - s)^{1 - \beta + \alpha}},$$

$s \in [0, t)$, which implies that $\| (-A)^{1-\beta+\alpha} T(t-s)g(s, (-A)^{-\alpha}y(s)) \|$ is integrable on $[0, t)$. We thus conclude that Ψ_2 is well defined and with values in $C([0, r] : X)$. It's clear from the previous remark that Ψ_1 is also well defined and with values in $C([0, r] : X)$.

It remain to show that the operator Ψ is a contraction on S . Firstly we prove that the range of Ψ is contained in S . Let y be a function in S . Then for $t \in [0, r]$ we get,

$$\begin{aligned} & \| \Psi(y)(t) - (-A)^\alpha x_0 \| \\ & \leq \| (T(t) - I)(-A)^\alpha (x_0 + g(0, x_0)) \| + \| (-A)^\alpha g(0, x_0) - (-A)^\alpha g(t, (-A)^{-\alpha}y(t)) \| \\ & \quad + \int_0^t \frac{C_{1-\beta+\alpha}}{(t-s)^{1-\beta+\alpha}} \| (-A)^\beta g(s, (-A)^{-\alpha}y(s)) \| ds \\ & \quad + \int_0^t \frac{C_\alpha}{(t-s)^\alpha} \| f(s, (-A)^{-\alpha}y(s)) \| ds \\ & \leq \frac{2(1-\mu)\delta}{6} + \| (-A)^{\alpha-\beta} \| L\{r^{\gamma_1} + \| (-A)^\alpha x_0 - y(t)\| \} \\ & \quad + \int_0^t \left(\frac{C_{1-\beta+\alpha}C_1}{(t-s)^{1-\beta+\alpha}} + \frac{C_\alpha C_1}{(t-s)^\alpha} \right) ds \\ & \leq \frac{2(1-\mu)\delta}{6} + \| (-A)^{\alpha-\beta} \| L\{r^{\gamma_1} + \delta\} + C_{1-\beta+\alpha}C_1 \frac{r^{\beta-\alpha}}{\beta-\alpha} + C_\alpha C_1 \frac{r^{1-\alpha}}{1-\alpha}. \end{aligned}$$

From the choice of r we conclude that

$$\| \Psi(y) - (-A)^\alpha x_0 \|_{r \leq \delta}$$

so that, $\Psi(y) \in S$.

On the other hand for $x(\cdot), y(\cdot) \in S$ and $t \in [0, r]$ we find

$$\begin{aligned} & \| \Psi(y)(t) - \Psi(x)(t) \| \\ & \leq \| (-A)^\alpha g(t, (-A)^{-\alpha}y(t)) - (-A)^\alpha g(t, (-A)^{-\alpha}x(t)) \| \\ & \quad + \int_0^t \frac{C_{1-\beta+\alpha}}{(t-s)^{1-\beta+\alpha}} \| (-A)^\beta g(s, (-A)^{-\alpha}y(s)) - (-A)^\beta g(s, (-A)^{-\alpha}x(s)) \| ds \\ & \quad + \int_0^t \frac{C_\alpha}{(t-s)^\alpha} \| f(s, (-A)^{-\alpha}y(s)) - f(s, (-A)^{-\alpha}x(s)) \| ds \\ & \leq \| (-A)^{\alpha-\beta} \| L \| y(t) - x(t) \| + \int_0^t \left\{ \frac{LC_{1-\beta+\alpha}}{(t-s)^{1-\beta+\alpha}} + \frac{NC_\alpha}{(t-s)^\alpha} \right\} \| y - x \|_r ds, \end{aligned}$$

thus

$$\| \Psi(y) - \Psi(x) \|_{r \leq} \left(\| (-A)^{\alpha-\beta} \| + LC_{1-\beta+\alpha} \frac{r^{\beta-\alpha}}{\beta-\alpha} + NC_\alpha \frac{r^{1-\alpha}}{1-\alpha} \right) \| y - x \|_r .$$

The last estimate and the choice of b imply that Ψ is a contraction mapping on S . Let $y(\cdot)$ be the unique fixed point of the operator Ψ in S . We affirm that $y(\cdot)$ is locally Hölder continuous. In fact, let ϑ be a real number with $0 < \vartheta < \min\{1 - \alpha, \beta - \alpha\}$ and $\vartheta + \beta > 1$, and let $\tilde{C} > 0$ be the constant guaranteed in lemma (1.1), such that for all $0 \leq s \leq t \leq T$ and $0 < h < 1$

$$\| (T(h) - I)(-A)^\alpha T(t-s) \| \leq \frac{\tilde{C}h^\vartheta}{(t-s)^{\vartheta+\alpha}}, \quad 0 \leq s < t$$

and

$$\| (T(h) - I)(-A)^{1-\beta+\alpha} T(t-s) \| \leq \frac{\tilde{C}h^\vartheta}{(t-s)^{1-\beta+\alpha+\vartheta}}, \quad 0 \leq s < t.$$

For $t \in [0, r)$ and $h > 0$ sufficiently small

$$\begin{aligned} & \| y(t+h) - y(t) \| \\ & \leq \| (T(h) - I)(-A)^\alpha T(t)(x_0 - g(0, x_0)) \| \\ & \quad + \| (-A)^{\alpha-\beta} \| L\{h^{\gamma_1} + \| y(t+h) - y(t) \| \} \\ & \quad + \int_0^t \| (T(h) - I)(-A)^{1-\beta+\alpha} T(t-s)(-A)^\beta g(s, (-A)^{-\alpha} y(s)) \| ds \\ & \quad + \int_t^{t+h} \| (-A)^{1-\beta+\alpha} T(t+h-s)(-A)^\beta g(s, (-A)^{-\alpha} y(s)) \| ds \\ & \quad + \int_0^t \| (T(h) - I)(-A)^\alpha T(t-s)f(s, (-A)^{-\alpha} y(s)) \| ds \\ & \quad + \int_t^{t+h} \| (-A)^\alpha T(t+h-s)f(s, (-A)^{-\alpha} y(s)) \| ds \\ & \leq \frac{\tilde{C}}{t^{\alpha+\vartheta}} \| x_0 - g(0, x_0) \| h^\vartheta + L \| (-A)^{\alpha-\beta} \| \{h^{\gamma_1} + \| y(t+h) - y(t) \| \} \\ & \quad + \int_0^t \frac{\tilde{C}h^\vartheta C_1}{(t-s)^{1-\beta+\alpha+\vartheta}} ds + \int_t^{t+h} \frac{C_{1-\beta+\alpha} C_1}{(t+h-s)^{1-\beta+\alpha}} ds \\ & \quad + \int_0^t \frac{\tilde{C}h^\vartheta C_1}{(t-s)^{\alpha+\vartheta}} ds + \int_t^{t+h} \frac{C_\alpha C_1}{(t+h-s)^\alpha} ds \\ & \leq \frac{C(x_0)h^\vartheta}{t^{\vartheta+\alpha}} + C_2 h^{\gamma_1} + L \| (-A)^{\alpha-\beta} \| \| y(t+h) - y(t) \| + C_3 h^\vartheta + C_4 h^{\beta-\alpha} + C_5 h^{1-\alpha} \end{aligned}$$

where the constants C_i are independent of t . If $\bar{\rho} = \min\{\vartheta, \gamma_1\}$, the last inequality can be rewritten in the form

$$\| y(t+h) - y(t) \| \leq \frac{C(\alpha, \beta, \vartheta, t, x_0)}{1 - \mu} h^{\bar{\rho}} \quad (13)$$

since $\mu = L \| (-A)^{\alpha-\beta} \| < 1$. Therefore the function $y(\cdot)$ is locally $\bar{\rho}$ -Hölder continuous on $(0, r)$, moreover, we can assume that $\bar{\rho} + \beta > 1$.

Now it is easy to prove that $s \rightarrow (-A)^\beta g(s, (-A)^{-\alpha} y(s))$ and $s \rightarrow f(s, (-A)^{-\alpha} y(s))$ are ρ -Hölder continuous on $(0, r)$, where $\rho = \min\{\bar{\rho}, \gamma_2\}$ and $\rho + \beta > 1$. From this remark, Theorem 2.4.1 in [5] and Lemma (2.1) below, we infer that the function

$$\begin{aligned} x(t) &= T(t)(x_0 + g(0, x_0)) - g(t, (-A)^{-\alpha} y(t)) \\ &\quad + \int_0^t (-A)^{1-\beta} T(t-s) (-A)^\beta g(s, (-A)^{-\alpha} y(s)) ds \\ &\quad + \int_0^t T(t-s) f(s, (-A)^{-\alpha} y(s)) ds \end{aligned} \quad (14)$$

is X_α -valued, that the integral terms in (14) are functions in $C^1([0, r] : X)$ and that $x(t) \in D(A)$ for all $t \in (0, r)$. Operating on $x(\cdot)$ with $(-A)^\alpha$, we conclude that $(-A)^{-\alpha} y = x$ and hence that $x(t) + g(t, x(t))$ is a C^1 function on $(0, b)$. The proof is completed.

The proof of the next Lemma is analogous to the proof of Theorem (2.4.1) in [5]. However there are some differences that require special attention and we include the principal ideas of this proof for completeness.

LEMMA 2.1. *Let $0 < \beta < 1$ and $g \in C([0, T] : X_{1-\beta})$. Assume that $g : [0, T] \rightarrow X$ is θ -Hölder continuous in $(0, T)$ with $\beta + \theta > 1$. If $y : [0, T] \rightarrow X$ is defined by*

$$y(t) = \int_0^t (-A)^{1-\beta} T(t-s) g(s) ds,$$

then $y(t) \in D(A)$ for every $t \in [0, T]$ and $\dot{y} \in C([0, T] : X)$.

Proof. For $t \in [0, T]$ we rewrite $y(t)$ in the form

$$\int_0^t (-A)^{1-\beta} T(t-s) (g(s) - g(t)) ds + \int_0^t (-A)^{1-\beta} T(t-s) g(t) ds = v(t) + w(t). \quad (15)$$

Clearly, $Aw(t) = T(t)(-A)^{1-\beta} g(t) - (-A)^{1-\beta} g(t) \in C([0, T] : X)$. For $\epsilon > 0$, sufficiently small we define the function

$$v_\epsilon(t) := \begin{cases} \int_0^{t-\epsilon} (-A)^{1-\beta} T(t-s) (g(s) - g(t)) ds, & \text{for } t \in [\epsilon, T], \\ 0 & \text{for } t \in [0, \epsilon]. \end{cases} \quad (16)$$

It is clear that $v_\epsilon(t) \in D(A)$. Moreover for $0 < \delta_1 < \delta_2$

$$\begin{aligned} \| Av_{\delta_2}(t) - Av_{\delta_1}(t) \| &\leq \int_{t-\delta_2}^{t-\delta_1} \| (-A)^{2-\beta} T(t-s) (g(s) - g(t)) \| ds \\ &\leq C_{2-\beta} (\delta_2^{\beta+\theta-1} - \delta_1^{\beta+\theta-1}). \end{aligned}$$

The last inequality proves that Av_δ is convergent, $\beta + \theta > 1$, and therefore

$$A(v(t)) = \int_0^t A^{2-\beta} T(t-s)(g(s) - g(t)) ds, \tag{17}$$

since A is a closed operator. Follows from the previous remark that $y(t) \in D(A)$ for $t \in [0, T]$. The continuity of $\partial_t y$ follows as in [5], Theorem 2.4.1.

In the rest of this paper for a function $j : [0, b] \times X \rightarrow X$ and $h \in \mathbf{R}$ we denote by $\partial_h j$ to the function

$$\partial_h j(t) = \frac{j(t+h) - j(t)}{h}.$$

Moreover, if j is differentiable we will employ the decomposition:

$$j(t+s, y) - j(t, y) = D_1 j(t, y) s + W_1(j, t, t+s, y) \tag{18}$$

and

$$j(t, y+y_1) - j(t, y) = D_2 j(t, y) \cdot y_1 + W_2(j, t, y, y+y_1) \tag{19}$$

where

$$\begin{aligned} \frac{W_1(j, t, t+s, y)}{|s|} &\rightarrow 0 & as & \quad s \rightarrow 0, \\ \frac{W_2(j, t, y, y+y_1)}{\|y_1\|} &\rightarrow 0 & as & \quad y_1 \rightarrow 0. \end{aligned}$$

In order to prove the next theorem, we need a preliminary result which is interesting in its own right.

LEMMA 2.2. *Assume that assumptions in Theorem (2.2) holds. If $x_0 \in D(A)$ and $g(0, x_0) \in D(A)$, then $x(\cdot) = x(\cdot, x_0)$ is Lipschitz on closed intervals.*

Proof. Initially we prove that $x(\cdot)$ is β -Hölder continuous on a closed interval $[0, b]$. Using the continuity of $(-A)^\beta g$ and f we can to assert that $(-A)^\beta g(s, x(s))$ and $f(s, x(s))$ are bounded by $C_1 > 0$ on $[0, b]$. Employing that $x_0 \in D(A)$ and that $g(0, x_0) \in D(A)$; for $t \in [0, b)$ and $h > 0$ we have

$$\| x(t+h) - x(t) \|$$

$$\begin{aligned}
&\leq C_2 h + \|g(t+h, x(t+h)) - g(t, x(t))\| \\
&\quad + \int_0^t \frac{C_{1-\beta}}{(t-s)^{1-\beta}} \|(-A)^\beta g(s+h, x(s+h)) - (-A)^\beta g(s, x(s))\| ds \\
&\quad + \int_0^h \|(-A)^{1-\beta} T(t+h-s) (-A)^\beta g(s, x(s))\| ds \\
&\quad + \tilde{M} \int_0^t \|f(s+h, x(s+h)) - f(s, x(s))\| ds + \tilde{M} \int_0^h \|f(s, x(s))\| ds \\
&\leq C_3 h + L \|(-A)^{-\beta}\| \|x(t+h) - x(t)\| \\
&\quad + \int_0^t \left\{ \frac{C_{1-\beta} L}{(t-s)^{1-\beta}} + N\tilde{M} \right\} \|x(s+h) - x(s)\| ds + \int_0^h \frac{C_{1-\beta}}{(t+h-s)^{1-\beta}} C_1 ds
\end{aligned}$$

thus

$$\begin{aligned}
\|x(t+h) - x(t)\| &\leq C_4 h^\beta + \|(-A)^{-\beta}\| L \|x(t+h) - x(t)\| \\
&\quad + \int_0^t \left\{ \frac{C_{1-\beta} L}{(t-s)^{1-\beta}} + N\tilde{M} \right\} \|x(s+h) - x(s)\| ds.
\end{aligned}$$

Since $\|(-A)^{-\beta}\| L < 1$, the Gronwall Bellman inequality (see Lemma 5.6.7 in [14]) implies that $x(\cdot)$ is β -Hölder continuous. Reiterating the previous estimates and using that $x(\cdot)$ is β -Hölder; if $t \in [0, T)$ and $h > 0$ we get

$$\|x(t+h) - x(t)\|$$

$$\begin{aligned}
&\leq C_5 h + \|(-A)^{-\beta}\| L \|x(t+h) - x(t)\| \\
&\quad + \int_0^t \left\{ \frac{C_{1-\beta} L}{(t-s)^{1-\beta}} + N\tilde{M} \right\} \|x(s+h) - x(s)\| ds \\
&\quad + \int_0^h \frac{C_{1-\beta} L}{(t+h-s)^{1-\beta}} \|(-A)^\beta g(s, x(s)) - (-A)^\beta g(0, x_0)\| ds \\
&\quad + \int_0^h \|T(t+h-s) (-A) g(0, x_0)\| ds
\end{aligned}$$

then

$$\begin{aligned}
\|x(t+h) - x(t)\| &\leq C_6 h^{2\beta} + \|(-A)^{-\beta}\| L \|x(t+h) - x(t)\| \\
&\quad + \int_0^t \left\{ \frac{C_{1-\beta} L}{(t-s)^{1-\beta}} + N\tilde{M} \right\} \|x(s+h) - x(s)\| ds.
\end{aligned}$$

The assumption $\|(-A)^{-\beta}\| L < 1$ and Gronwall Bellman inequality, implies that $x(\cdot)$ is 2β -Hölder continuous. Clearly the previous routine permit to infer that $x(\cdot)$ is Lipschitz continuous, this completes the proof.

In the next theorem we establish the existence of classical solutions using some usual regularity assumptions on the functions f and $(-A)^\beta g$.

THEOREM 2.4. *Assume that $(-A)^{1-\beta}g(\cdot)$ and $f(\cdot)$ are continuously differentiable functions on $[0, T] \times \Omega$. If $x_0, g(0, x_0) \in D(A)$ and $\| D_2g(0, x_0) \|_{\mathcal{L}(X)} < 1$ then $\dot{x}(\cdot, x_0) \in C([0, b] : X)$ for some $0 < b < T$.*

Proof: Let $x(\cdot) = x(\cdot, x_0)$ and $z(\cdot)$ a solution of the integral equation

$$\begin{aligned} z(t) &= T(t)(Ax_0 + Ag(0, x_0) + f(0, x_0)) + h(t) - [D_2g(t, x(t))](z(t)) \\ &\quad + \int_0^t (-A)^{1-\beta}T(t-s)[D_2(-A)^\beta g(s, x(s))](z(s))ds \\ &\quad + \int_0^t T(t-s)[D_2f(s, x(s))](z(s))ds \end{aligned} \tag{20}$$

where

$$z(0) = Ax_0 + Ag(0, x_0) + f(0, x_0) - D_1g(0, x_0) - [D_2g(0, x_0)](z(0))$$

and

$$\begin{aligned} h(t) &= -D_1g(t, x(t)) + \int_0^t (-A)^{1-\beta}T(t-s)D_1(-A)^\beta g(s, x(s))ds \\ &\quad + \int_0^t T(t-s)D_1f(s, x(s))ds. \end{aligned}$$

The existence and uniqueness of local solution for the integral equation (20) is consequence of the contraction mapping principle and the condition $\| D_2g(0, x_0) \|_{\mathcal{L}(X)} < 1$; we omit details. In what follows we assume that $x(\cdot)$ and $z(\cdot)$ are defined on $[0, 2b]$ where $0 < 2b < T$ and $\| D_2g(\theta, x_\theta) \|_{2b} < \eta < 1$.

Next we prove that $\dot{x} = z$ on $[0, b]$. To this end, we observe that from Lemma (2.2)

$$\frac{W_2((-A)^\beta g, s, x(s), x(s+h))}{\| x(s+h) - x(s) \|} \cdot \frac{\| x(s+h) - x(s) \|}{h} \rightarrow 0 \quad \text{as } h \rightarrow 0$$

and

$$\frac{W_2(f, s, x(s), x(s+h))}{\| x(s+h) - x(s) \|} \cdot \frac{\| x(s+h) - x(s) \|}{h} \rightarrow 0 \quad \text{as } h \rightarrow 0.$$

uniformly for $s \in [0, b]$. Consequently, for $t \in [0, b]$ and $h > 0$ we find

$$\| \xi(t, h) \| = \left\| \frac{x(t+h) - x(t)}{h} - z(t) \right\|$$

$$\begin{aligned}
&\leq \| T(t)(\frac{T(h) - I}{h}x_0 - A(x_0)) \| + \| \frac{1}{h} \int_0^h T(t+h-s)f(s, x(s))ds - T(t)f(0, x_0) \| \\
&+ \| T(t)(\frac{T(h) - I}{h})g(0, x_0) + \frac{1}{h} \int_0^h (-A)^{1-\beta}T(t+h-s)(-A)^\beta g(s, x(s))ds \| \\
&+ \| D_1g(t, x(t+h)) - D_1g(t, x(t)) \| + \| [D_2g(t, x(t))](\xi(t, h)) \| \\
&+ \| \frac{W_1(g, t, t+h, x(t+h))}{h} \| + \| \frac{W_2(g, t, x(t), x(t+h))}{h} \| \\
&+ \int_0^t \frac{C_{1-\beta}}{(t-s)^{1-\beta}} \| D_1(-A)^\beta g(s, x(s+h)) - D_1(-A)^\beta g(s, x(s)) \| ds \\
&+ \int_0^t \frac{C_{1-\beta}}{(t-s)^{1-\beta}} \| [D_2(-A)^\beta g(s, x(s))](\xi(s, h)) \| ds \\
&+ \int_0^t \frac{C_{1-\beta}}{(t-s)^{1-\beta}} \| \frac{W_1((-A)^\beta g, s, s+h, x(s+h))}{h} \| ds \\
&+ \int_0^t \frac{C_{1-\beta}}{(t-s)^{1-\beta}} \| \frac{W_2((-A)^\beta g, s, x(s), x(s+h))}{h} \| ds \\
&+ \int_0^t \tilde{M} \| D_1f(s, x(s+h)) - D_1f(s, x(s)) \| ds \\
&+ \int_0^t \tilde{M} \| [D_2f(s, x(s))](\xi(s, h)) \| ds + \tilde{M} \int_0^t \| \frac{W_1(f, s, s+h, x(s+h))}{h} \| ds \\
&+ \tilde{M} \int_0^t \| \frac{W_2(f, s, x(s), x(s+h))}{h} \| ds. \\
&\leq \rho(t, h) + \frac{1}{h} \int_0^h \| (-A)^{1-\beta}T(t+h-s)((-A)^\beta g(0, x_0) - (-A)^\beta g(s, x(s))) \| ds \\
&+ \| D_2g(t, x(t)) \| \| \xi(t, h) \| + \int_0^t \frac{C_{1-\beta}}{(t-s)^{1-\beta}} \| D_2(-A)^\beta g(s, x(s)) \| \| \xi(s, h) \| ds \\
&+ \tilde{M} \int_0^t \| D_2f(s, x(s)) \| \| \xi(s, h) \| ds
\end{aligned}$$

where $\rho(t, h) \rightarrow 0$ as $h \rightarrow 0$, uniformly for $t \in [0, b]$. Using that $\| D_2g(\cdot, x(\cdot)) \|_b < \eta$ and the Lipschitz continuity of $x(\cdot)$, follow that

$$\begin{aligned}
\| \xi(t, h) \| &\leq \frac{1}{1-\eta} \rho(t, h) + \frac{C_{1-\beta}LCh^\beta}{\beta} \\
&+ \frac{1}{1-\eta} \int_0^t \frac{C_{1-\beta}}{(t-s)^{1-\beta}} \| D_2(-A)^\beta g(s, x(s)) \| \| \xi(s, h) \| ds \\
&+ \frac{1}{1-\eta} \tilde{M} \int_0^t \| D_2f(s, x(s)) \| \| \xi(s, h) \| ds.
\end{aligned}$$

Finally, the Gronwall Bellman inequality (see Lemma 5.6.7 in [14]) show that $\xi(t, h) \rightarrow 0$ as $h \rightarrow 0$. Therefore $\dot{x}(\cdot, x_0) = z(\cdot)$. This completes the proof.

COROLLARY 2.1. *If g is a $D(A)$ -valued continuous function then there exists a unique classical solutions of (1) defined on $[0, b]$ for some $0 < b < T$.*

Proof: From Theorem (2.4) we know that $x(\cdot) = x(\cdot, x_0) \in C^1([0, b] : X)$ for some $0 < b < T$. Since $x(\cdot, x_0)$ is Lipschitz continuous in $[0, b]$, from Theorem 2.4.1 in [5] and Lemma (2.1) we infer that $x(t) + g(t, x(t)) \in D(A)$ for $t \in [0, b]$ and therefore that $x(t) \in D(A)$ for $t \in [0, b]$. The proof is complete.

3. EXAMPLES

In this section we sketch briefly some applications.

3.1. Functional Differential Equations with Unbounded Delay.

The regularity results obtained in this work are used to prove the existence of regular solutions, "Classical" and "N-Classical" solutions, for a class of quasi-linear neutral functional differential equations with unbounded delay that can be modeled in the form

$$\frac{d}{dt}(x(t) + F(t, x_t)) = Ax(t) + G(t, x_t), \quad t \geq \sigma, \quad (21)$$

$$x_\sigma = \varphi \in \mathcal{B}, \quad (22)$$

where A is the infinitesimal generator of an analytic semigroup of bounded linear operators $(T(t))_{t \geq 0}$ on a Banach space X ; the history $x_t : (-\infty, 0] \rightarrow X$, $x_t(\theta) = x(t + \theta)$, belongs to some abstract phase space \mathcal{B} defined axiomatically, as in Hale and Kato [3], and where the axioms are established employing the terminology and notations used in Hino-Murakami-Naito [9]. A complete reference including results of existence of mild, strong and periodical solutions for (21)-(22) are the papers [6], [7]. The existence of "N-Classical" and "Classical" solutions is studied in [8], actually in preparation.

3.2. Partial Differential Equations of Sobolev Type

There is an extensive literature on semi-linear Sobolev evolution equations modeled in the form

$$\frac{d}{dt}(Bu(t)) = Au(t) + f(t, u(t)), \quad t > 0, \quad (23)$$

$$u(0) = u_0, \quad u_0 \in D(B). \quad (24)$$

where A, B be closed linear operators on a Banach space X . The literature includes different and complete results concerning to existence, uniqueness and qualitative properties of mild, strong and classical solutions for (23)-(24) (see [1], [12], [16], [15]). Some usual assumptions on the operators A, B (see for example [1] and [12]) are

- a) A, B are closed linear operators.
 b) $D(B) \subset D(A)$ and B has a continuous inverse.

From a), b) and the Closed Graph Theorem it follows that AB^{-1} is a bounded linear operator on X . In this case the approach is to consider the related integral equation

$$x(t) = T(t)Bx_0 + \int_0^t T(t-s)f(s, B^{-1}x(s))ds,$$

where $T(t)$, $t \geq 0$, is the semigroup generated by AB^{-1} .

We shall consider the abstract Cauchy problem

$$\frac{d}{dt}(u(t) + Bu(t)) = Au(t) + f(t, u(t)), \quad t > 0, \quad (25)$$

$$u(0) = u_0, \quad u_0 \in D(B), \quad (26)$$

where A, B are closed linear operators on a Banach space X and

- a') $D(A) \subset D(B)$ and B has a continuous inverse
 b') AB^{-1} is the infinitesimal generator of an analytic semigroup of bounded linear operators on X .

Using the conditions a'), b'), we consider the associated system

$$\frac{d}{dt}(u + B^{-1}u(t)) = AB^{-1}u(t) + f(t, B^{-1}u(t)), \quad t > 0, \quad (27)$$

$$u(0) = B^{-1}u_0, \quad u_0 \in D(B). \quad (28)$$

If in addition, B^{-1} is $D(AB^{-1})$ -valued continuous and f is continuously differentiable, the existence of classical solutions for (27)-(28) and consequently for (25)-(26), follows from Corollary (2.1).

REFERENCES

1. Brill, Heinz "A semilinear Sobolev evolution equation in a Banach space". J. Differential Equations 24 (1977), no. 3, 412-425.
2. J. Dieudone, "Fundation in Modern Analysis", vol 1, Academic, New York 1969.
3. Hale, Jack K.; Kato, Junji. "Phase space for retarded equations with infinite delay". Funkcial. Ekvac. 21 (1978), no. 1, 11-41.
4. Hale, Jack K.; Verduyn Lunel, Sjoerd M. "Introduction to functional-differential equations". Applied Mathematical Sciences, 99. Springer-Verlag, New York, 1993.
5. Goldstein, Jerome A. "Semigroups of linear operators and applications". Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 1985.
6. Hernández, Eduardo; Henríquez, Hernán R. "Existence of periodic solutions of partial neutral functional-differential equations with unbounded delay". J. Math. Anal. Appl. 221 (1998), no. 2, 499-522.

7. Hernández, Eduardo; Henríquez, Hernán R. "Existence results for partial neutral functional-differential equations with unbounded delay". *J. Math. Anal. Appl.* 221 (1998), no. 2, 452–475.
8. Hernández, Eduardo; "Regular Solutions for Partial Neutral Functional Differential Equations with Unbounded Delay". Preprint.
9. Hino, Yoshiyuki; Murakami, Satoru; Yoshizawa, Taro. "Existence of almost periodic solutions of some functional-differential equations with infinite delay in a Banach space". *Tôhoku Math. J. (2)* 49 (1997), no. 1, 133–147.
10. Krein, S. G. "Linear differential equations in Banach space". Translated from the Russian by J. M. Danskin. *Translations of Mathematical Monographs*, Vol. 29. American Mathematical Society, Providence, R.I., 1971.
11. Lagnese, John E. "General boundary value problems for differential equations of Sobolev type". *SIAM J. Math. Anal.* 3 (1972), 105–119.
12. Lightbourne, James H., III; Rankin, Samuel M., III "A partial functional-differential equation of Sobolev type". *J. Math. Anal. Appl.* 93 (1983), no. 2, 328–337.
13. L. Nachbin, "Introduction to functional Analysis: Banach Space and Differential Calculus". Marcel Dekker, New York, 1981.
14. Pazy, A. "Semigroups of linear operators and applications to partial differential equations". *Applied Mathematical Sciences*, 44. Springer-Verlag, New York-Berlin, 1983.
15. Showalter, R. E. "Degenerate parabolic initial-boundary value problems". *J. Differential Equations* 31 (1979), no. 3, 296–312.
16. Showalter, R. E. "A nonlinear parabolic-Sobolev equation". *J. Math. Anal. Appl.* 50 (1975), 183–190.