

UNIVERSIDADE DE SÃO PAULO  
Instituto de Ciências Matemáticas e de Computação  
ISSN 0103-2577

---

**Bilipschitz Determinacy of Quasihomogeneous Germs**

Alexandre Cesar Gurgel Fernandes  
Maria Aparecida Soares Ruas

N<sup>o</sup> 139

---

NOTAS

Série Matemática



São Carlos – SP  
Abr./2002

SYSNO	<u>1253541</u>
DATA	<u>1</u> / <u>1</u>
ICMC - SBAB	

## Bilipschitz determinacy of quasihomogeneous germs

Alexandre Cesar Gurgel Fernandes\*

*Centro de Ciências*  
*Universidade Federal do Ceará*  
*Av. Humberto Monte, s/n Campus do Pici - Bloco 914*  
*Fortaleza, Ceará, Brazil*  
*e*  
*Depto. de Matemática*  
*ICMC/USP- São Carlos*  
*Caixa Postal 668*  
*13560-970 São Carlos SP, Brazil*  
E-mail: alex@icmc.sc.usp.br

and

Maria Aparecida Soares Ruas†

*Depto. de Matemática*  
*ICMC/USP- São Carlos*  
*Caixa Postal 668*  
*13560-970 São Carlos SP, Brazil*  
E-mail: maasruas@icmc.sc.usp.br

### Abstract

We obtain estimates for the degree of bilipschitz determinacy of quasihomogeneous function-germs.

### Resumo

Neste trabalho, obtemos estimativas para o grau de determinação bilipschitz de germes de funções quase-homogêneas.

\* Research partially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico, (CNPq), Brazil, grant # 140499/00-8.

† Research partially supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico, (CNPq), Brazil, grant # 300066/88-0, and by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Brazil, grant # 97/10735-3.

## 1. INTRODUCTION

A basic problem in Singularity Theory is the local classification of mappings module diffeomorphisms. In 1965, H. Whitney justified the rigidity of the classification problem by  $C^1$ -diffeomorphism giving the following example:

$$F_t(x, y) = xy(x - y)(x - ty); \quad 0 < t < 1 \quad (1)$$

which presents the following phenomenon: for any  $t \neq s$  in  $I = (0, 1)$  it is not possible to construct a  $C^1$ -diffeomorphism  $\phi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$  such that  $F_t = F_s \circ \phi$ . This motivated the classification of mappings by "isomorphisms" weaker than diffeomorphisms.

There is an extensive literature related to  $C^r$ -equivalence ( $1 \leq r < \infty$ ) of map-germs, among them [5], [4] and [1] which are more closely related to this work. However, only few recent works deal with the problem of bilipschitz classification of map-germs. This work is inspired in a recent paper by J.-P. Henry and A. Parusinski [2], where they show that the bilipschitz equivalence of analytic function-germs admits continuous moduli. We obtain estimates for the degree of bilipschitz determinacy of quasihomogeneous function-germs. Examples are given to show that the estimates are sharp.

## 2. BILIPSCHITZ EQUIVALENCE

Let  $\lambda \in \mathbb{R}$  be a positive number. A mapping  $\phi : U \subset \mathbb{R}^n \rightarrow \mathbb{R}^p$  is called  $\lambda$ -Lipschitz, or simply Lipschitz if it satisfies:

$$\|\phi(x) - \phi(y)\| \leq \lambda \|x - y\| \quad \forall x, y \in U.$$

When  $n = p$  and  $\phi$  has a Lipschitz inverse, we say that  $\phi$  is bilipschitz.

Two germs  $f, g : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^p, 0)$  are called bilipschitz equivalent if there exists a bilipschitz map-germ  $\phi : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^n, 0)$  such that  $f = g \circ \phi$ .

EXAMPLE 2.2.1. Let  $f, g : (\mathbb{R}, 0) \rightarrow (\mathbb{R}, 0)$  be given by  $f(x) = x, g(x) = x^3$ . It is easy to show that  $f$  and  $g$  are not bilipschitz equivalent. On the other hand, there is a homeomorphism  $\phi : (\mathbb{R}, 0) \rightarrow (\mathbb{R}, 0)$  such that  $f = \phi \circ g$ .

Let  $f : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}, 0)$  be the germ of an analytic function,

$$f(x) = f_m(x) + f_{m+1}(x) + \dots,$$

with  $f_i$  a homogeneous form of degree  $i$ , and  $f_m \neq 0$ . We denote by  $m_f := m$ , the multiplicity of  $f$ . We say that  $f$  has non-degenerate tangent cone if  $0 \in \mathbb{R}^n$  is the only point in  $\mathbb{R}^n$  in which

$$\frac{\partial f_m}{\partial x_1} = \dots = \frac{\partial f_m}{\partial x_n} = 0.$$

PROPOSITION 2.2.2. Let  $f : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}, 0)$  be the germ of an analytic function. Then

$$m_f = \text{ord}_r[\text{sup}|f|_{B(0,r)}],$$

where  $B(0, r)$  denote the ball centered at the origin with radius  $r$ .

*Proof.* Let  $\alpha = \text{ord}_r[\text{sup}|f|_{B(0,r)}]$ . Write

$$f(x) = f_m(x) + f_{m+1}(x) + \cdots$$

with  $f_i$  a homogeneous form of degree  $i$ , and  $f_m \neq 0$ . Let  $x = (x_1, \dots, x_n)$  be such that  $f_m(x) \neq 0$ . Then, given  $r > 0$  we have

$$\begin{aligned} |f(rx)| &= r^m |f_m(x) + rf_{m+1}(x) + \cdots| \\ &\geq Kr^m \end{aligned}$$

for some constant  $K > 0$ , hence  $m \geq \alpha$ .

On the other hand, from the Curve Selection Lemma, there exists an analytic arc  $\gamma : [0, \epsilon) \rightarrow \mathbb{R}^n$ ,  $\gamma(0) = 0$ , such that

$$\alpha = \text{ord}_r |f(\gamma(r))|$$

and  $|\gamma(r)| \leq r$  for each  $r > 0$ . Since  $\gamma(0) = 0$ , we can write  $\gamma(r) = r\tilde{\gamma}(r)$  with  $\lim_{r \rightarrow 0} \tilde{\gamma}(r) < \infty$ . Therefore,

$$\begin{aligned} |f(\gamma(r))| &= r^m |f_m(\tilde{\gamma}(r)) + rf_{m+1}(\tilde{\gamma}(r)) + \cdots| \\ &\leq Lr^m \end{aligned}$$

for some constant  $L > 0$ . Hence,  $m \leq \alpha$ . ▀

**COROLLARY 2.2.3.** *Let  $f, g : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}, 0)$  be germs of analytic functions. If  $f$  and  $g$  are bilipschitz equivalent, then  $m_f = m_g$ .*

The corollary above in the complex case was proved by J.-J. Risler and D. Trotman in [3]. It is obvious that the converse statement is false, but we can prove the following result

**PROPOSITION 2.2.4.** *Let  $F_t : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}, 0)$  be a smooth family of smooth function-germs. If  $m_{F_t}$  is constant and  $F_t$  has non-degenerate tangent cone for each  $t$ , then for each  $t \neq s$ ,  $F_t$  and  $F_s$  are bilipschitz equivalent.*

The result above will follow as consequence of Theorem 3.3.3.

**COROLLARY 2.2.5.** *The family (1) satisfies:  $F_t$  and  $F_s$  are bilipschitz equivalent  $\forall t, s \in (0, 1)$ .*

It is valuable to observe that the Proposition 2.2.4 does not guarantee the non-rigidity of the bilipschitz classification problem for analytic functions. In fact, J.-P. Henry and A. Parusinski ([2]) presented the family  $F_t : (\mathbb{C}^2, 0) \rightarrow (\mathbb{C}^2, 0)$  given by  $F_t(x, y) = x^3 - 3t^2xy^2 + y^6$  which satisfies: for any  $t \neq s \in (0, \frac{1}{2})$  there is no bilipschitz map-germ  $\phi : (\mathbb{C}^2, 0) \rightarrow (\mathbb{C}^2, 0)$  such that  $F_t = F_s \circ \phi$ . The proof is based on the analysis of the expansion of the germs of the family along each arc of their polar curves. The argument in [2] also holds in the real case, that is, the following holds:

**PROPOSITION 2.2.6.** *The family  $F_t : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$  given by  $F_t(x, y) = x^3 - 3t^2xy^2 + y^6$  satisfies: for any  $t \neq s \in (0, \frac{1}{2})$  there is no bilipschitz map  $\phi : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}^2, 0)$  such that  $F_t = F_s \circ \phi$ .*

Note that  $F_t$  is a deformation of the quasihomogeneous germ  $f = x^3 + y^6$  which has an isolated singularity at origin. Therefore, it is natural to ask for which  $\theta(x, y)$  the family  $f + t\theta$  is bilipschitz trivial.

### 3. BILIPSCHITZ DETERMINACY OF QUASIHOMOGENEOUS GERMS

Let  $f_t : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}, 0)$   $t \in I$  (an interval in  $\mathbb{R}$ ), be a smooth family of smooth function-germs. That is, there is a neighborhood  $U$  of 0 in  $\mathbb{R}^n$  and a smooth function  $F : U \times I \rightarrow \mathbb{R}$  such that  $F(0, t) = 0$  and  $f_t(x) = F(x, t) \forall t \in I, \forall x \in U$ . We call  $f_t$  *strongly bilipschitz trivial* when there is a continuous family of  $\lambda$ -Lipschitz map-germs (vector field)  $v_t : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^n, 0)$  such that

$$\frac{\partial f_t}{\partial t}(x) = (df_t)_x(v_t(x))$$

$\forall t \in \mathbb{R}$  and  $\forall x$  near 0 in  $\mathbb{R}^n$ .

**THEOREM 3.3.1.** *If  $f_t$  is bilipschitz trivial, then for each  $t \neq s \in I$  there is a bilipschitz map-germ  $\phi : (\mathbb{R}^n, 0) \rightarrow (\mathbb{R}^n, 0)$  such that  $f_t = f_s \circ \phi$ .*

The theorem above is known as a result of Thom-Levine type and its proof is immediate, since the flow of a Lipschitz vector field is bilipschitz.

Let  $\mathcal{E}_n$  be the space of smooth function-germs  $(\mathbb{R}^n, 0) \rightarrow \mathbb{R}$ . Given  $f \in \mathcal{E}_n$ , we denote  $Nf(x) = \sum \left[ \frac{\partial f}{\partial x_i}(x) \right]^2$ . We say that  $Nf(x)$  satisfies a *Lojasiewicz condition* if there exist constants  $c > 0$  and  $\alpha > 0$  such that  $Nf(x) \geq c\|x\|^\alpha$ .

Fix the weights  $(r_1, \dots, r_n)$ . We recall that a function  $f$  is called *quasihomogeneous* with respect to the given weights if there is a number  $d$  such that  $f$  satisfies the following equation:

$$f(\lambda \cdot x) = \lambda^d f(x_1, \dots, x_n)$$

$\forall \lambda \in \mathbb{R}$  and  $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ , where  $\lambda \cdot x = (\lambda^{r_1} x_1, \dots, \lambda^{r_n} x_n)$ . With respect to the given weights, for each monomial  $x^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$ , we define  $\text{fil}(x^\alpha) = \sum_{i=1}^n \alpha_i r_i$ . We define a filtration in the ring  $\mathcal{E}_n$  via the function  $\text{fil}(f) = \min\{\text{fil}(x^\alpha) : (\frac{\partial f}{\partial x^\alpha})(0) \neq 0\}$ , for each  $f \in \mathcal{E}_n$ . We can extend this definition to  $\mathcal{E}_{n+1}$ , the ring of 1-parameter families of smooth function-germs in  $\mathcal{E}_n$ , by defining  $\text{fil}(x^\alpha t^\beta) = \text{fil}(x^\alpha)$ .

Let  $(r_1, \dots, r_n; 2k)$  be fixed. The standard control function of type  $(r_1, \dots, r_n; 2k)$  is  $\rho(x) = x^{2\alpha_1} + \cdots + x^{2\alpha_n}$ , where the  $\alpha_i$  are chosen such that the function  $\rho$  is quasihomogeneous of type  $(r_1, \dots, r_n; 2k)$ .

**LEMMA 3.3.2.** *Let  $h(x)$  be a quasihomogeneous polynomial function of type  $(r_1, \dots, r_n; 2k)$ , with  $r_1 \leq \cdots \leq r_n$ ,  $\rho$  the standard control function of same type that  $h$  and  $h_t(x)$  a deformation of  $h$  such that:*

$$\text{fil}(h_t) \geq 2k + r_n, \quad t \in [0, 1]. \quad (2)$$

*Then the function  $\frac{h_t(x)}{\rho(x)}$  is  $c$ -Lipschitz, with  $c$  independent of  $t$ .*

*Proof.* Without loss of generality, we can suppose that  $h_t(x)$  is quasihomogeneous of type  $(r_1, \dots, r_n; d)$  where  $d \geq 2k + r_n$ . We consider  $G_t(x, y) : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  given by  $G_t(x, y) = |\rho(y)h_t(x) - \rho(x)h_t(y)|$ ,  $m_t(x, y) = \|x - y\|\rho(x)\rho(y)$  and  $M = \{(x, y, t) : G_t(x, y) = 1\}$ . Since  $M$  is closed, the number  $c = \inf\{m_t(x, y) : (x, y, t) \in M\}$  is positive. Now, let  $x, y \in \mathbb{R}^n$  be sufficiently near the origin  $x \neq 0, y \neq 0$  and  $x \neq y$ . Let  $\lambda > 0$  be such that  $G_t(\lambda \cdot x, \lambda \cdot y) = 1$ , that is,

$$G_t(x, y) = \frac{1}{\lambda^{2k+d}} \quad (3)$$

On the other hand, we use that  $\lambda > 1$  to obtain:

$$\begin{aligned} m_t(\lambda \cdot x, \lambda \cdot y) &= \lambda^{4k} \|\lambda \cdot x - \lambda \cdot y\| \rho(x) \rho(y) \\ &\leq \lambda^{4k+r_n} \|x - y\| \rho(x) \rho(y) \\ &= \lambda^{4k+r_n} m_t(x, y) \end{aligned}$$

$\therefore$

$$m_t(x, y) \geq \frac{1}{\lambda^{4k+r_n}} c. \quad (4)$$

Now, we use that  $\lambda > 1$ ,  $d \geq 2k + r_n$ , (3), (3) and we obtain the following inequality  $m_t(x, y) \geq cG_t(x, y)$ , that is,

$$\left\| \frac{h_t(x)}{\rho(x)} - \frac{h_t(y)}{\rho(y)} \right\| \leq c^{-1} \|x - y\|.$$

**■**

**THEOREM 3.3.3.** *Let  $f : \mathbb{R}^n, 0 \rightarrow \mathbb{R}^n, 0$  be the germ of a quasihomogeneous polynomial function of type  $(r_1, \dots, r_n; d)$ ,  $r_1 \leq \dots \leq r_n$  with isolated singularity. Let  $f_t(x) = f(x) + t\Theta(x, t)$ ,  $t \in [0, 1]$ , be a smooth deformation of  $f$ . If  $\text{fil}(\Theta) \geq d + r_n - r_1$ , then  $f_t$  admits a strong bilipschitz trivialization along  $I = [0, 1]$ .*

*Proof.* We can see that for each  $i$  there exists a  $s_i$  such that  $\frac{\partial f}{\partial x_i}$  is quasihomogeneous of the type  $(r_1, \dots, r_n; s_i)$ ,  $s_i = d - r_i$ .

Let  $N^*f$  be defined by

$$N^*f = \sum \left[ \frac{\partial f}{\partial x_i} \right]^{2\alpha_i},$$

where  $\alpha_i = \frac{k}{s_i}$  and  $k = \text{l.c.m.}(s_i)$ . Therefore  $N^*f$  is a quasihomogeneous control function of the type  $(r_1, \dots, r_n; 2k)$ .

The lemma bellow is proved in [4].

**LEMMA 3.3.4.** *There exist constants  $0 < c_2 < c_1$  such that*

$$c_2\rho(x) \leq N^*f_t(x) \leq c_1\rho(x).$$

We have the following equation;

$$\frac{\partial f_t}{\partial t} N^*f_t = df_t(W),$$

where  $W$  is given by

$$W = \sum W_i \frac{\partial}{\partial x_i} \text{ where } W_i = \frac{\partial f_t}{\partial t} \left[ \frac{\partial f}{\partial x_i} \right]^{2\alpha_i - 1}.$$

Since  $\text{fil} \left( \frac{\partial f_t}{\partial t} \right) \geq d + r_n - r_1$  and

$$\begin{aligned} \text{fil} \left( \left[ \frac{\partial f_t}{\partial x_i} \right]^{2\alpha_i - 1} \right) &= (2\alpha_i - 1) \text{fil} \left( \frac{\partial f_t}{\partial x_i} \right) \\ &= (2\alpha_i - 1)(d - r_i) \\ &= 2K - d + r_i \\ &\geq 2k - d + r_1 \end{aligned}$$

we have that  $\min \text{fil}(W_i) \geq \text{fil}(\Theta) + 2k - d + r_1 \geq 2k + r_n$ .

Let  $v : \mathbb{R}^n \times \mathbb{R}, 0 \rightarrow \mathbb{R}^n \times \mathbb{R}, 0$  be the vector field given by  $\frac{W}{N^*f_t}$ . From Lemma 3.3.2, it follows that  $v$  is a Lipschitz vector field.

Finally the equation  $(\frac{\partial f_t}{\partial t})(x) = (df_t)_x(v(x, t))$  gives the strong bilipschitz triviality of the family  $f_t(x)$  along a small open interval around  $t = 0$ . Since the same argument is true for each  $t \in I$ , the proof is complete.  $\blacksquare$

The following result shows that the estimate given in Theorem 3.3.3 is sharp.

PROPOSITION 3.3.5. *Let  $f_t : (\mathbb{R}^2, 0) \rightarrow (\mathbb{R}, 0)$ ;  $t \in I = (-\delta, \delta) \subset \mathbb{R}$  be given by*

$$f_t(x, y) = \frac{1}{3}x^3 - t^2xy^{3n-2} + y^{3n}.$$

*Then  $f_t$  is not strongly bilipschitz trivial.*

REMARK 3.3.6. Let  $f(x, y) = \frac{1}{3}x^3 + y^{3n}$ . Note that  $f$  is quasihomogeneous of type  $(n, 1; 3n)$ . From Theorem 3.3.3 it follows that  $f + t\theta$  is strongly bilipschitz trivial for each  $\theta(x, t)$  such that  $fil(\theta) \geq 4n - 1$ .

*Proof (of the Proposition 3.3.5).* Let  $m = 3n - 2$ . Here we repeat the argument-proof from Theorem 1.1 in [2]. Suppose that  $v(x, y, t) = v_1(x, y, t)\frac{\partial}{\partial x} + v_2(x, y, t)\frac{\partial}{\partial y}$  is a vector field such that:

$$\left(\frac{\partial f_t}{\partial t}\right)(x, y) = (df_t)_x(v(x, y, t))$$

The polar curve of  $f_t$   $\{(x, y) \in \mathbb{R}^2 : \frac{\partial f_t}{\partial x}(x, y) = 0\}$  is equal to the set  $\{(x, y) \in \mathbb{R}^2 : x^2 = t^2y^m\}$ . Thus,  $v_1(ty^{m/2}, y, t)$  and  $v_2(-ty^{m/2}, y, t)$  satisfy:

$$v_1(ty^{m/2}, y, t)\frac{\partial f_t}{\partial y}(ty^{m/2}, y, t) = -\frac{\partial f_t}{\partial t}(ty^{m/2}, y, t) \quad (5)$$

$$v_2(-ty^{m/2}, y, t)\frac{\partial f_t}{\partial y}(-ty^{m/2}, y, t) = -\frac{\partial f_t}{\partial t}(-ty^{m/2}, y, t). \quad (6)$$

From equations (5) and (6) we have:

$$v_1(ty^{m/2}, y, t) = \frac{2t^2y^{m/2-1}}{-mt^3y^{m/2-2} + 3n}$$

$$v_2(-ty^{m/2}, y, t) = \frac{-2t^2y^{m/2-1}}{mt^3y^{m/2-2} + 3n}$$

Thus,

$$v_1(ty^{m/2}, y, t) - v_2(-ty^{m/2}, y, t) \sim y^{m/2-1} \quad (7)$$



On the other hand,

$$\|(ty^{m/2}, y, t) - (-ty^{m/2}, y, t)\| \sim y^{m/2} \quad (8)$$

But, (7) and (8) show that  $v_2$  is not Lipschitz. Hence  $f$  is not strongly bilipschitz trivial. ■

The invariant for bilipschitz equivalence  $\text{Inv}(f_t)$  presented in [2] is independent of  $t$ , hence does not distinguish the element of the given family  $f_t$ .

### REFERENCES

1. S. Bromberg and S. L. de Medrano,  *$C^r$ -sufficiency of quasihomogeneous functions*. Real and Complex Singularities, Pitman Research Notes in Mathematics Series 333, (1995), 179–189.
2. A. Parusinski and J.-P. Henry, *Existence of moduli for bilipschitz equivalence of analytic functions*. Prépublications de l'Université d'Angers no. 137, 2001.
3. J.-J. Risler and D. Trotman, *bilipschitz invariance of the multiplicity*. Bull. London Math. Soc. 29 (1997), no. 2, 200–204.
4. M.A.S. Ruas and M.J. Saia,  *$C^l$ -determinacy of weighted homogeneous germs*. Hokkaido Mathematical Journal, vol. 26, no 1 (1997), 89–99.
5. C. T. C. Wall, *Finite determinacy of smooth map-germs*. Bull. London Math. Society 13 (1981), 418–539.

# NOTAS DO ICMC

## SÉRIE MATEMÁTICA

- 138/2002 MANCINI, S.; MANOEL, M.; TEIXEIRA, M.A. – Divergent diagrams of folds associated with a set of involutions.
- 137/2002 LÓPEZ, A. – A structure theorem for foliations on non-compact two-manifolds.
- 136/2002 COHN, P.M.; FERREIRA, V.O. – Split firs.
- 135/2002 HERNÁNDEZ M., E., HENRÍQUEZ, H. – Global solutions for a functional second order abstract Cauchy problem with nonlocal conditions.
- 134/2002 HERNÁNDEZ M., E., HENRÍQUEZ, H. – Existence of solutions of the functional second order abstract Cauchy problem with nonlocal conditions.
- 133/2002 HERNÁNDEZ M., E. – A massera type criterion for partial neutral functional differential equations.
- 132/2002 FERREIRA, V. O.; GONÇALVES, J.Z.; MANDEL, A. – Free symmetric and unitary pairs in division rings with involutions.
- 131/2001 CARBINATTO, M.C.; RYBAKOWSKI, K.P. – Morse decompositions in the absence of uniqueness.
- 130/2001 CONDE, A. – O terceiro problema de Hilbert.
- 129/2001 SAIA, M.J.; TOMAZELLA, J.N. – Deformations with constant Milnor number and multiplicity of non-degenerate complex hypersurfaces.