

UNIVERSIDADE DE SÃO PAULO  
Instituto de Ciências Matemáticas e de Computação  
ISSN 0103-2577

---

On Codimensions  $k$  Immersions of  $m$ -Manifolds for  
 $k = m - 3$ ,  $k = m - 5$  and  $k = m - 6$

Carlos Biasi  
Alice K. M. Libardi  
Denise de Mattos  
Edivaldo L. dos Santos

Nº 309

---

NOTAS

Série Matemática



São Carlos – SP  
Abr./2009

SYSNO	1739978
DATA	/ /
ICMC - SBAB	

On codimensions  $k$  immersions of  $m$ -manifolds for  $k = m - 3$ ,  $k = m - 5$  and  
 $k = m - 6$

Carlos Biasi \*

*Departamento de Matemática, Instituto de Ciências Matemáticas e de Computação, Universidade de São Paulo - Campus de São Carlos, Caixa Postal 668, 13560-970 São Carlos SP, Brazil*  
E-mail: biasi@icmc.usp.br

Alice K. M. Libardi†

*IGCE-UNESP, 13500-230, Rio Claro, SP, Brazil*  
E-mail: alicekml@rc.unesp.br

Denise de Mattos‡

*Departamento de Matemática, Instituto de Ciências Matemáticas e de Computação, Universidade de São Paulo - Campus de São Carlos, Caixa Postal 668, 13560-970 São Carlos SP, Brazil*  
E-mail: deniseml@icmc.usp.br

Edivaldo L. dos Santos§

*Departamento de Matemática, Universidade Federal de São Carlos-UFSCar, Caixa Postal 676, 13565-905, São Carlos-SP, Brazil*  
E-mail: edivaldo@dm.ufscar.br

Considere  $M$  uma variedade suave, fechada, conexa de dimensão  $n$ ,  $N$  uma variedade suave de dimensão  $n$  e  $f : M \rightarrow N$  uma função contínua com codimensão  $k = n - m > 0$ . Neste artigo, sob certas condições, nós provamos que  $f$  é homotópica à uma imersão, nos seguintes casos:  $m \equiv 1(4)$  e codimensão  $k = m - 3$ ;  $m \equiv 2(4)$ ,  $m \equiv 3(8)$  e codimensão  $k = m - 5$ ;  $m \equiv 2(4)$ ,  $m \equiv 3(8)$  e codimensão  $k = m - 6$ . Este trabalho complementa resultados de Biasi et al. (Manus. Math. 126, 527-530, 2008 e Manus. Math. 104, 97-110, 2001); Koschorke (Lecture Notes in Mathematics, vol. 1350. Springer, Heidelberg, 1988); e de Bang-He Li e Gui-Song Li (Math. Proc. Camb. Phil. Soc. 112, 281-285, 1992).

\* Partially supported by FAPESP 06/03932-8, Brazil.

† Partially supported by FAPESP 06/03932-8, Brazil.

‡ Partially supported by FAPESP 06/03932-8 and CNPq 308390/2008-3, Brazil.

§ Partially supported by FAPESP 06/03932-8 and CNPq 304480/2008-8, Brazil.

Let us consider  $M$  a closed smooth connected  $m$ -manifold,  $N$  a smooth  $n$ -manifold and  $f : M \rightarrow N$  a continuous map with codimension  $k = n - m > 0$ . In this paper, under certain conditions, we prove that  $f$  is homotopic to an immersion, in the following cases:  $m \equiv 1(4)$  and codimension  $k = m - 3$ ;  $m \equiv 2(4)$ ,  $m \equiv 3(8)$  and codimension  $k = m - 5$ ;  $m \equiv 2(4)$ ,  $m \equiv 3(8)$  and codimension  $k = m - 6$ . This work complements some results of Biasi et al. (Manus. Math. 126, 527-530, 2008 and Manus. Math. 104, 97-110, 2001); Koschorke (Lecture Notes in Mathematics, vol. 1350. Springer, Heidelberg, 1988); and of Bang-He Li and Gui-Song Li (Math. Proc. Camb. Phil. Soc. 112, 281-285, 1992).

## 1. INTRODUCTION

Let  $M$  be a smooth  $m$ -manifold,  $N$  be a smooth  $n$ -manifold, and let  $f : M \rightarrow N$  be a continuous map with codimension  $k = n - m > 0$ . If  $k = m - 1$  it was proved in [6] that  $f$  is always homotopic to an immersion. If  $k = m - 2$ , in general,  $f$  is not homotopic to an immersion (see [7]). For this case, recently C. Biasi and A. K. M. Libardi in [1] proved that, on certain conditions,  $f$  is homotopic to an immersion. More specifically, Theorem 2.1 of [1] states that: “Let  $M$  be a closed smooth connected manifold of dimension  $m \equiv 1(4)$ , and let  $N$  be a smooth manifold of dimension  $2m - 2$ . Let  $f : M \rightarrow N$  be a continuous map such that  $f_* : H_1(M; \mathbb{Z}_2) \rightarrow \check{H}_1(f(M); \mathbb{Z}_2)$  is injective. Then  $f$  is homotopic to an immersion.”

In this paper, we prove a version of [1, Theorem 2.1] on existence of immersions for codimensions  $k = m - 3$ ,  $k = m - 5$  and  $k = m - 6$ . Specifically, we prove the following theorem:

**THEOREM 1.1.** *Let  $M$  be a closed smooth connected manifold of dimension  $m$ ,  $N$  be a smooth manifold of dimension  $2m - i$  and let  $f : M \rightarrow N$  be a continuous map. Let us suppose that  $f_* : H_{i-1}(M; \mathbb{Z}_2) \rightarrow \check{H}_{i-1}(f(M); \mathbb{Z}_2)$  is injective and that one of the following conditions is satisfied:*

- (a)  $i = 3$  and  $m \equiv 1(4)$ ,  
 $H_1(M; \mathbb{Z}_2) = 0$  and  $H^{m-2}(M; \mathbb{Z})$  does not have elements of order 4.
- (b)  $i = 5$  and  $m \equiv 2(4)$ ,  
 $H_1(M; \mathbb{Z}_2) = H_2(M; \mathbb{Z}_8) = H_3(M; \mathbb{Z}_2) = 0$ .
- (c)  $i = 5$  and  $m \equiv 3(8)$ ,  
 $H_3(M; \mathbb{Z}_4) = 0$ ,  $H_1(M; \mathbb{Z}_2) = H_1(M; \mathbb{Z}_{12}) = 0$  (or  $H_1(M; \mathbb{Z}) = 0$ ) and  $H^{m-4}(M; \mathbb{Z})$  does not have elements of order 4.
- (d)  $i = 6$  and  $m \equiv 2(4)$ ,  
 $H_1(M; \mathbb{Z}_2) = H_2(M; \mathbb{Z}) = H_3(M; \mathbb{Z}_2) = H_4(M; \mathbb{Z}_2) = 0$  and  $H^{m-5}(M; \mathbb{Z})$  does not have elements of order 4.

- (e)  $i = 5$  and  $m \equiv 3(8)$ ,  
 $f^*w_1(N) = w_1(M)$ ,  $H_2(M; \mathbb{Z}_2) = H_3(M; \mathbb{Z}_8) = H_4(M; \mathbb{Z}_2) = 0$ .

Then  $f$  is homotopic to an immersion.

We observe that this theorem complement some results of [1, 2], [4, 5] and [9], and that this result cover all possible cases which could be obtained by using the Pachter's table [11].

## 2. PRELIMINARES AND NOTATIONS

Let  $M$  be a closed connected smooth manifold of dimension  $m$ ,  $N$  be a smooth manifold of dimension  $n$ , and  $f : M \rightarrow N$  a continuous map with codimension  $k = n - m > 0$ . Let  $[M] \in H_m(M; \mathbb{Z}_2)$  denote the fundamental classe of  $M$  and  $U_f \in H^k(N; \mathbb{Z}_2)$  the Poincaré dual of  $f_*[M] \in H_n^C(N; \mathbb{Z}_2)$ , i.e.,  $f_*[M] = D(U_f) = U_f \frown [N]$ , where  $[N] \in H_n^C(N; \mathbb{Z}_2)$  is the fundamental class of  $N$  and  $D$  is the Poincaré dual isomorphism.

We define a homology class  $\theta(f) \in H^{m-k}(M; \mathbb{Z}_2)$  by

$$\theta(f) = D(f^*(U_f) - w_k(f)) \in H_{m-k}(M; \mathbb{Z}_2),$$

where  $w_k(f) = w_k(\nu_f)$  is the  $k^{\text{th}}$  Stiefel-Whitney class of the stable normal bundle of  $f$  (see [2] for more details). We recall that  $\theta(f)$  denotes the primary obstruction for  $f$  be homotopic to an embedding, that is, if  $f$  is homotopic to an embedding then  $\theta(f) = 0$ .

Throughout the paper,  $\check{H}$  denotes the Čech singular homology group,  $\pi_{m,q}^p$  denotes the  $(m+p)^{\text{th}}$  homotopy group of the Stiefel manifold  $V_{m+q,q}$ , i.e.,  $\pi_{m,q}^p = \pi_{m+p}(V_{m+q,q})$  and  $\rho$  denotes the reduction modulo two homomorphism.

## 3. PROOF OF THE THEOREM 1.1

In this section, let us observe that all obstruction classes lie in cohomological groups with local coefficients system, however, under our assumptions the local coefficients system is always trivial.

*Proof* ( Proof of Theorem 1.1 (a)). Let us consider  $f_1 : M \rightarrow N \times \mathbb{R}$  defined by  $f_1(x) = (f(x), 0)$ , for each  $x \in M$ . Since  $H_1(M; \mathbb{Z}_2) = 0$  by our assumption, we have that  $f_{1*} : H_1(M; \mathbb{Z}_2) \rightarrow \check{H}_1(f_1(M); \mathbb{Z}_2)$  is an injective map. Also  $m \equiv 1(4)$  and  $N \times \mathbb{R}$  has dimension  $2m - 2$ . Therefore, it follows from [1, Theorem 2.1] that  $f_1$  is homotopic to an immersion  $g_1$ .

On the other hand, Lemmas 6.1 and 6.2 of [2] imply that  $f_1^*(U_{f_1}) = 0 \in H^{m-2}(M; \mathbb{Z}_2)$ . If we consider  $\tilde{f} : M \rightarrow f_1(M)$ , defined by  $\tilde{f}(x) = f_1(x)$ ,  $x \in M$ , by [3, Theorem 1.1], we have that  $\tilde{f}_*(\theta(f_1)) = 0$ , where  $\tilde{f}_* : H_2(M; \mathbb{Z}_2) \rightarrow \check{H}_2(f_1(M); \mathbb{Z}_2)$ . Since  $\tilde{f}_*$  is injective by assumption, we have that  $\tilde{f}_*$  is injective. Therefore,  $0 = \theta(f_1) = D(f_1^*(U_{f_1}) - w_{m-2}(f_1))$  and  $w_{m-2}(f_1) = w_{m-2}(g_1) = w_{m-2}(f) = 0$ .

Let us consider the stable normal bundle  $\nu_{g_1}^{m-2}$ . Since the Stiefel manifold  $V_{m-2,1}$  is  $(m-4)$ -connected,  $\nu_{g_1}^{m-2}$  has an 1-frame field over the  $(m-3)$ -skeleton of  $M$ . The



obstruction to the existence of a 1-frame field over the  $(m-2)$ -skeleton of  $M$  is the class  $W_{m-2}(g_1) \in H^{m-2}(M; \pi_{m-3}(V_{m-2,1}))$ . Since  $m-3 = 4s+2$  it follows from [11] that  $\pi_{m-3}(V_{m-2,1}) = \pi_{m-3,1}^0 = \pi_{4s+2,1}^0 = \mathbb{Z}$ . By our assumption,  $H^{m-2}(M; \mathbb{Z})$  does not have elements of order 4 and since  $\rho(W_{m-2}(g_1)) = w_{m-2}(g_1) = 0$ , where  $\rho : H^{m-2}(M; \pi_{m-3}(V_{m-2,1})) \rightarrow H^{m-2}(M; \mathbb{Z}_2)$ , we have that  $W_{m-2}(g_1)$  vanishes. Then there exists a 1-frame field over the  $(m-2)$ -skeleton of  $M$ . The obstruction to the existence of a 1-frame field over the  $(m-1)$ -skeleton of  $M$  is the class  $W_{m-1}(g_1) \in H^{m-1}(M; \pi_{m-2}(V_{m-2,1})) = H^{m-1}(M; \pi_{m-3,1}^1)$  and  $\pi_{m-3,1}^1 = \mathbb{Z}_2$  (see [11]). By hypothesis,  $H_1(M; \mathbb{Z}_2) = 0$ , then  $W_{m-1}(g_1) \in H^{m-1}(M; \pi_{m-2}(V_{m-2,1}))$  vanishes and there exists a 1-frame field over the  $(m-1)$ -skeleton of  $M$ .

Now, let us consider  $g_q : M \rightarrow (N \times \mathbb{R}) \times \mathbb{R}^{q-1}$  defined by  $g_q(x) = (g_1(x), 0)$ ,  $q \geq 4$ . Since  $\nu_{g_1}^{m-2}$  has a 1-frame field over the  $(m-1)$ -skeleton of  $M$ , we have that  $\nu_{g_q}^{m-3+q}$  has a  $q$ -frame field over the  $(m-1)$ -skeleton of  $M$  and the obstruction to the existence of a  $q$ -frame field over the  $m$ -skeleton of  $M$  is the class  $W_m(g_q) \in H^m(M; \pi_{m-1}(V_{m-3+q,q})) = H^m(M; \pi_{m-3,q}^2)$ . Since  $\pi_{m-3,q}^2 = 0$ , for  $q \geq 4$  (see [11]), we have that the obstruction class  $W_m(g_q) \in H^m(M; \pi_{m-1}(V_{m-3+q,q}))$  vanishes. Hence by a result of Hirsch [8], we conclude that  $f$  is homotopic to an immersion.  $\blacksquare$

*Proof* (Proof of Theorem 1.1 (b)-(c)). In this proof we consider simultaneously the cases  $m \equiv 2(4)$  and  $m \equiv 3(8)$ . Let us consider  $f_1 : M \rightarrow N \times \mathbb{R}$  defined by  $f_1(x) = (f(x), 0)$ , for each  $x \in M$ . By Lemmas 6.1 and 6.2 of [2] we see that  $f_1^*(U_{f_1}) = 0 \in H^{m-4}(M; \mathbb{Z}_2)$ . If we consider  $\tilde{f} : M \rightarrow f_1(M)$ , defined by  $\tilde{f}(x) = f_1(x)$ ,  $x \in M$ , by [3, Theorem 1.1], we have that  $\tilde{f}_*(\theta(f_1)) = 0$ , where  $\tilde{f}_* : H_4(M; \mathbb{Z}_2) \rightarrow H_4(f_1(M); \mathbb{Z}_2)$ . Since  $f_*$  is injective by our assumption, we have that  $\tilde{f}_*$  is injective. Therefore  $0 = \theta(f_1) = D(f_1^*(U_{f_1}) - w_{m-4}(f_1))$  and  $w_{m-4}(f_1) = w_{m-4}(f) = 0$ .

Let us consider  $f_q : M \rightarrow N \times \mathbb{R}^q$ ,  $q \geq 6$ . Since  $N \times \mathbb{R}^q$  has dimension  $2m-5+q \geq 2m+1$  we have that  $f_q$  is homotopic to an immersion  $g_q$ . Therefore,  $w_{m-4}(g_q) = w_{m-4}(f_q) = w_{m-4}(f_1) = w_{m-4}(f) = 0$ .

Let us consider the stable normal bundle  $\nu_{g_q}^{m-5+q}$ . Since the Stiefel manifold  $V_{m-5+q,q}$  is  $(m-6)$ -connected,  $\nu_{g_q}^{m-5+q}$  has a  $q$ -frame field over the  $(m-5)$ -skeleton of  $M$ . The obstruction to the existence of a  $q$ -frame field over the  $(m-4)$ -skeleton of  $M$  is the class  $W_{m-4}(g_q) \in H^{m-4}(M; \pi_{m-5}(V_{m-5+q,q}))$ . Since  $m-5 = 4s+1$  or  $m-5 = 4s+2$  and  $q \geq 6$  it follows from [11] that  $\pi_{m-5}(V_{m-5+q,q}) = \pi_{m-5,q}^0 = \pi_{4s+1,q}^0 = \mathbb{Z}_2$  or  $\pi_{m-5}(V_{m-5+q,q}) = \pi_{m-5,q}^0 = \pi_{4s+2,q}^0 = \mathbb{Z}$ . In the first case, the homomorphism  $\rho : H^{m-4}(M; \pi_{m-5}(V_{m-5+q,q})) \rightarrow H^{m-4}(M; \mathbb{Z}_2)$  is an isomorphism and  $\rho(W_{m-4}(g_q)) = w_{m-4}(g_q) = 0$ . In the second case,  $H^{m-4}(M; \mathbb{Z})$  does not have elements of order 4 by our assumption and  $\rho(W_{m-4}(g_q)) = w_{m-4}(g_q) = 0$ . In both cases, the obstruction class  $W_{m-4}(g_q)$  vanishes and then there exists a  $q$ -frame field over the  $(m-4)$ -skeleton of  $M$ .

The obstruction class to the existence of a  $q$ -frame field over the  $(m-3)$ -skeleton of  $M$  is the class  $W_{m-3}(g_q) \in H^{m-3}(M; \pi_{m-4}(V_{m-5+q,q}))$ , which vanishes, because the homotopy group  $\pi_{m-4}(V_{m-5+q,q}) = \pi_{m-5,q}^1 = \pi_{4s+1,q}^1 = \mathbb{Z}_2$  or  $\pi_{m-4}(V_{m-5+q,q}) = \pi_{m-5,q}^1 = \pi_{4s+2,q}^1 = \mathbb{Z}_4$ , for  $q \geq 6$  (see [11]) and by hypothesis,  $H_3(M; \mathbb{Z}_2) = 0$  or  $H_3(M; \mathbb{Z}_4) = 0$ .

The next obstruction class is  $W_{m-2}(g_q) \in H^{m-2}(M; \pi_{m-3}(V_{m-5+q,q}))$ . Since the homotopy group  $\pi_{m-3}(V_{m-5+q,q}) = \pi_{m-5,q}^2 = \pi_{4s+1,q}^2 = \mathbb{Z}_8$  or  $\pi_{m-3}(V_{m-5+q,q}) = \pi_{m-5,q}^2 =$

$\pi_{4s+2,q}^2 = 0$ , for  $q \geq 6$  and by assumption  $H_2(M; \mathbb{Z}_8) = 0$ , we have that the obstruction class  $W_{m-2}(g_q)$  vanishes and there exists a  $q$ -frame field over the  $(m-2)$ -skeleton of  $M$ .

The next obstruction class is  $W_{m-1}(g_q) \in H^{m-1}(M; \pi_{m-2}(V_{m-5+q,q}))$ . Since the homotopy group  $\pi_{m-2}(V_{m-5+q,q}) = \pi_{m-5,q}^3 = \pi_{4s+1,q}^3 = \mathbb{Z}_2$  or  $\pi_{m-2}(V_{m-5+q,q}) = \pi_{m-5,q}^3 = \pi_{4s+2,q}^3 = \mathbb{Z}_{12}$ , for  $q \geq 6$  (see [11]) and by assumption  $H_1(M; \mathbb{Z}_2) = 0$  or  $H_1(M; \mathbb{Z}_{12}) = 0$ , we have that the obstruction class  $W_{m-1}(g_q)$  vanishes and there exists a  $q$ -frame field over the  $(m-1)$ -skeleton of  $M$ .

Finally, the obstruction to the existence of a  $q$ -frame field over the  $m$ -skeleton of  $M$  is the class  $W_m(g_q) \in H^m(M; \pi_{m-1}(V_{m-5+q,q}))$ . Since  $m-5 = 4s+5$  or  $m-5 = 8s-2$  and  $q \geq 6$  it follows from [11] that  $\pi_{m-1}(V_{m-5+q,q}) = \pi_{m-5,q}^4 = \pi_{4s+5,q}^4 = 0$  or  $\pi_{m-1}(V_{m-5+q,q}) = \pi_{m-5,q}^4 = \pi_{8s-2,q}^4 = 0$ , and consequently,  $W_m(g_q)$  vanishes. Therefore, by Hirsch [8], we conclude that  $f$  is homotopic to an immersion.  $\blacksquare$

*Proof* ( Proof of Theorem 1.1(d)). Let us consider  $f_1: M \rightarrow N \times \mathbb{R}$  defined by  $f_1(x) = (f(x), 0)$ , for each  $x \in M$ . Since  $H_4(M; \mathbb{Z}_2) = 0$ , we have that  $f_{1*}: H_4(M; \mathbb{Z}_2) \rightarrow \check{H}_4(f_1(M); \mathbb{Z}_2)$  is an injective map. Also  $m \equiv 2(4)$ ,  $H_1(M; \mathbb{Z}_2) = H_2(M; \mathbb{Z}) = H_3(M; \mathbb{Z}_2) = 0$  and  $N \times \mathbb{R}$  has dimension  $2m-5$ . By Theorem 1.1(b), we see that  $f_1$  is homotopic to an immersion  $g_1$ .

On the other hand, since  $f_*: H_5(M; \mathbb{Z}_2) \rightarrow \check{H}_5(f(M); \mathbb{Z}_2)$  is injective, by similar argument to that in the proof of Theorem 1.1(b)-(c), we have that  $0 = \theta(f_1) = D(f_1^*(U_{f_1}) - w_{m-5}(f_1))$  and  $w_{m-5}(f_1) = w_{m-5}(f) = 0$ .

Let us consider the stable normal bundle  $\nu_{g_1}^{m-5}$ . Since  $V_{m-5,1}$  is  $(m-7)$ -connected,  $\nu_{g_1}^{m-5}$  has an 1-frame field over the  $(m-6)$ -skeleton of  $M$ . Therefore, the obstruction to the existence of an 1-frame field over the  $(m-5)$ -skeleton of  $M$  is the class  $W_{m-5}(g_1) \in H^{m-5}(M; \pi_{m-6}(V_{m-5,1}))$ . Since  $m-6 = 4s$  it follows from [11] that  $\pi_{m-6}(V_{m-5,1}) = \pi_{m-6,1}^0 = \pi_{4s,1}^0 = \mathbb{Z}$ . Since  $H^{m-5}(M; \mathbb{Z})$  does not have elements of order 4 and  $\rho(W_{m-5}(g_1)) = w_{m-5}(g_1) = 0$ , where  $\rho: H^{m-5}(M; \pi_{m-6}(V_{m-5,1})) \rightarrow H^{m-5}(M; \mathbb{Z}_2)$ , we have that  $W_{m-5}(g_1)$  vanishes. Then, there exists an 1-frame field over the  $(m-5)$ -skeleton of  $M$ .

The next obstruction class is  $W_{m-4}(g_1) \in H^{m-4}(M; \pi_{m-5}(V_{m-5,1}))$ . Since the homotopy group  $\pi_{m-5}(V_{m-5,1}) = \pi_{m-6,1}^1 = \pi_{4s,1}^1 = \mathbb{Z}_2$  (see [11]) and by hypothesis,  $H_4(M; \mathbb{Z}_2) = 0$ , we see that the obstruction class  $W_{m-4}(g_1)$  vanishes. Thus, there exists a 1-frame field over the  $(m-4)$ -skeleton of  $M$ .

The obstruction to the existence of an 1-frame field over the  $(m-3)$ -skeleton of  $M$  is the class  $W_{m-3}(g_1) \in H^{m-3}(M; \pi_{m-4}(V_{m-5,1}))$ , which vanishes, because  $\pi_{m-4}(V_{m-5,1}) = \pi_{m-6,1}^2 = \pi_{4s,1}^2 = \mathbb{Z}_2$  and by hypothesis  $H_3(M; \mathbb{Z}_2) = 0$ .

The next obstruction class is  $W_{m-2}(g_1) \in H^{m-2}(M; \pi_{m-3}(V_{m-5,1}))$ . Since  $m-6 = 4s$ , we have that  $m-6 = 8s$  or  $m-6 = 8s+4$ . Therefore,  $\pi_{m-3}(V_{m-5,1}) = \pi_{m-6,1}^3 = \mathbb{Z}_{24}$  (see [11]) and by assumption  $H_2(M; \mathbb{Z}_{24}) = 0$ , we have that the obstruction class  $W_{m-2}(g_1)$  vanishes. Thus, there exists an 1-frame field over the  $(m-2)$ -skeleton of  $M$ .

The obstruction to the existence of a 1-frame field over the  $(m-1)$ -skeleton of  $M$  is the class  $W_{m-1}(g_1) \in H^{m-1}(M; \pi_{m-2}(V_{m-5,1}))$ . Since  $\pi_{m-2}(V_{m-5,1}) = \pi_{m-6,1}^4 = \pi_{4s,1}^4 = 0$ , we see that  $W_{m-1}(g_1)$  vanishes. Thus, there exists an 1-frame field over the  $(m-1)$ -skeleton of  $M$ .



Finally, the obstruction to the existence of an 1-frame field over the  $m$ -skeleton of  $M$  is the class  $W_m(g_1) \in H^m(M; \pi_{m-1}(V_{m-5,1}))$ , which vanishes, since  $\pi_{m-1}(V_{m-5,1}) = \pi^5_{m-6,1} = \pi^5_{4s,1} = 0$ . Then, by Hirsch [8], we conclude that  $f$  is homotopic to an immersion.  $\blacksquare$

*Proof* ( Proof of Theorem 1.1(e)). Let us consider  $f_1 : M \rightarrow N \times \mathbb{R}$  defined by  $f_1(x) = (f(x), 0)$ , for each  $x \in M$ . Since  $f_* : H_5(M; \mathbb{Z}_2) \rightarrow \check{H}_5(f(M); \mathbb{Z}_2)$  is injective, by similar argument to that in the proof of Theorem 1.1(b)-(c), we have that  $0 = \theta(f_1) = D(f_1^*(U_{f_1}) - w_{m-5}(f_1))$  and  $w_{m-5}(f_1) = w_{m-5}(f) = 0$ .

Also, if we consider  $f_q : M \rightarrow N \times \mathbb{R}^q$ ,  $q \geq 7$ ,  $N \times \mathbb{R}^q$  has dimension  $2m - 6 + q \geq 2m + 1$ . Therefore  $f_q$  is homotopic to an immersion  $g_q$  and  $w_{m-5}(g_q) = w_{m-5}(f_q) = w_{m-5}(f_1) = w_{m-5}(f) = 0$ .

Let us consider the stable normal bundle  $\nu_{g_q}^{m-6+q}$ . Since the Stiefel manifold  $V_{m-6+q,q}$  is  $(m-7)$ -connected,  $\nu_{g_q}^{m-6+q}$  has a  $q$ -frame field over the  $(m-6)$ -skeleton of  $M$ . The obstruction to the existence of a  $q$ -frame field over the  $(m-5)$ -skeleton of  $M$  is the class  $W_{m-5}(g_q) \in H^{m-5}(M; \pi_{m-6}(V_{m-6+q,q}))$ . Since  $m-6 = 4s+1$  it follows from [11] that  $\pi_{m-6}(V_{m-6+q,q}) = \pi_{m-6,q}^0 = \pi_{4s+1,q}^0 = \mathbb{Z}_2$ . In this case, the homomorphism  $\rho : H^{m-5}(M; \pi_{m-6}(V_{m-6+q,q})) \rightarrow \check{H}^{m-5}(M; \mathbb{Z}_2)$  is an isomorphism and  $\rho(W_{m-5}(g_q)) = w_{m-5}(g_q) = 0$ . Therefore,  $W_{m-5}(g_q)$  vanishes and there exists a  $q$ -frame field over the  $(m-5)$ -skeleton of  $M$ .

The next obstruction class is  $W_{m-4}(g_q) \in H^{m-4}(M; \pi_{m-5}(V_{m-6+q,q}))$ . Since the homotopy group  $\pi_{m-5}(V_{m-6+q,q}) = \pi_{m-6,q}^1 = \pi_{4s+1,q}^1 = \mathbb{Z}_2$ , for  $q \geq 7$  (see [11]) and by hypothesis,  $H_4(M; \mathbb{Z}_2) = 0$ , we have that the obstruction class  $W_{m-4}(g_q)$  vanishes. Thus there exists a  $q$ -frame field over the  $(m-4)$ -skeleton of  $M$ .

The obstruction to the existence of a  $q$ -frame field over the  $(m-3)$ -skeleton of  $M$  is the class  $W_{m-3}(g_q) \in H^{m-3}(M; \pi_{m-4}(V_{m-6+q,q}))$ , which vanishes, because  $\pi_{m-4}(V_{m-6+q,q}) = \pi_{m-6,q}^2 = \pi_{4s+1,q}^2 = \mathbb{Z}_8$ , for  $q \geq 7$  and by hypothesis  $H_3(M; \mathbb{Z}_8) = 0$ .

The next obstruction class is  $W_{m-2}(g_q) \in H^{m-2}(M; \pi_{m-3}(V_{m-6+q,q}))$ . Since the homotopy group  $\pi_{m-3}(V_{m-6+q,q}) = \pi_{m-6,q}^3 = \pi_{4s+1,q}^3 = \mathbb{Z}_2$ , for  $q \geq 7$  (see [11]) and by assumption  $H_2(M; \mathbb{Z}_2) = 0$ , we have that the obstruction class  $W_{m-2}(g_q)$  vanishes and there exists a  $q$ -frame field over the  $(m-2)$ -skeleton of  $M$ .

The next obstruction class is  $W_{m-1}(g_q) \in H^{m-1}(M; \pi_{m-2}(V_{m-6+q,q}))$ . Since  $m-6 = 4s+5$  it follows from [11] that  $\pi_{m-2}(V_{m-6+q,q}) = \pi_{m-6,q}^4 = \pi_{4s+5,q}^4 = 0$ , and consequently,  $W_{m-1}(g_q)$  vanishes. Thus, there exists a  $q$ -frame field over the  $(m-1)$ -skeleton of  $M$ .

Finally, the obstruction to the existence of a  $q$ -frame field over the  $m$ -skeleton of  $M$  is the class  $W_m(g_q) \in H^m(M; \pi_{m-1}(V_{m-6+q,q}))$ . Since  $m-6 = 8s+5$  and  $q \geq 7$  it follows from [11] that  $\pi_{m-1}(V_{m-6+q,q}) = \pi_{m-6,q}^5 = \pi_{8s+5,q}^5 = 0$ , and consequently,  $W_m(g_q)$  vanishes. Then, by Hirsch [8], we have that  $f$  is homotopic to an immersion.  $\blacksquare$

## REFERENCES

1. Biasi, C. and Libardi, A.K.M. *On codimensions  $k$  immersions of  $m$ -manifolds for  $k = 1$  and  $k = m - 2$* , Manus. Math. **126**, 527-530 (2008)

2. Biasi, C., Daccach J. and Saeki, O. *A primary obstruction to topological embeddings and its applications*, Manus. Math. **104**, 97-110 (2001)
3. Biasi, C. and Saeki, O. *On the self intersection set and the image of a generic map*, Math. Scand. **80**, 5-24 (1997)
4. Li, B.H. *Existence of immersions of  $n$ -manifolds in manifolds of dimension  $2m - 2$  or  $2m - 3$* . J. Sys. Sci. & Math. Sci **6** (3), 177-185 (1986)
5. Li, B.H. and Li, G.S. *Immersions with non-zero normal vector fields*. Math. Proc. Camb. Phil. Soc. **112**, 281-285 (1992)
6. Li, B.H. and Peterson, F.P. *On immersions of  $k$ -manifolds in  $(2k - 1)$ - manifolds*, Proc. Amer. Math. Soc. **83**, n. 1, 159-162 (1981)
7. Li, B.H. and Peterson, F.P. *On immersions of  $n$ -manifolds in  $(2n - 2)$ - manifolds*, Proc. Amer. Math. Soc. **97**, n. 3, 531-538 (1986)
8. M.W. Hirsch, *Immersions of manifolds*. Trans. Amer. Math. Soc. **93**, 242-276 (1959)
9. U. Koschorke, *The singularity method and immersions of  $m$ -manifolds into manifolds of dimensions  $2m - 2$ ,  $2m - 3$  and  $2m - 4$* . Lecture Notes in Mathematics, **1350** - Springer Verlag (1988)
10. J. Milnor and J. Stasheff, *Characteristic Classes*. Ann. of Math. Studies, vol. 76 Princeton Univ. Press, Princeton, N.J. (1974)
11. G.F. Paechter, *The groups  $\pi_r(V_{n,m})$* , Quart. J. Math. Oxford(2), **7**, 249-268 (1956)
12. E.H. Spanier, *Algebraic Topology*, McGraw-Hill Book Co., New York (1966)



# NOTAS DO ICMC

## SÉRIE MATEMÁTICA

- 308/09 BARBOSA, G.F.; GRULHA JR., N.; SAIA, M.J. – Stable types and the minimal Whitney stratification of discriminants.
- 307/09 ARRIETA, J.M.; CARVALHO, A.N.; LANGA, J.A.; RODRIGUEZ-BERNAL, A. – Continuity of dynamical structures for non-autonomous evolution equations under singular perturbations.
- 306/09 CARVALHO, A.N.; LANGA, J.A.; ROBINSON, J.C. – Finite-dimensional global attractors in banach spaces.
- 305/09 CARABALLO, T.; CARVALHO, A.N.; LANGA, J.A.; RIVERO, F. – Existence of pullback attractors for pullback asymptotically compact processes.
- 304/09 CARABALLO, T.; LANGA, J.A.; RIVERO, F.; CARVALHO, A.N. – A gradient-like non-autonomous evolution process.
- 303/09 CAMPOS, J.E.P.P. – Distinguishing link-homotopy classes by algebraic methods.
- 302/08 BUOSI, M.; IZUMIYA, S.; RUAS, M.A. – Horo-tight circles in hyperbolic space.
- 301/08 ARRIETA, J.M.; CARVALHO, A.N.; LOZADA-CRUZ, G. – Dynamics in dumbbell domains III. Continuity of attractors.
- 300/08 ARRIETA, J.M.; CARVALHO, A.N.; LOZADA-CRUZ, G. – Dynamics in dumbbell domains II. The limiting problem.
- 299/08 DEMUNER, D.P.; FEDERSON, M.; GUTIERREZ, C. – The Poincaré-Bendixson theorem on the Klein bottle for continuous vector fields.